

resolution of PET to define the precise locations of  $^{18}\text{F}$ -NaF uptake in coronary arteries.

Finally, we do agree with Dr. Mohler and colleagues that, after further validation,  $^{18}\text{F}$ -NaF PET of the coronary arteries may give unique insights into the pathophysiology of calcium deposition and perhaps its dispersal with appropriate therapy.

**\*Marc R. Dweck, MD**  
**Nikhil V. Joshi, MD**  
**Nicholas A. Boon, MD**  
**James H. F. Rudd, PhD**  
**David E. Newby, DSc**

\*Centre for Cardiovascular Science  
 University of Edinburgh  
 Little France Crescent  
 Edinburgh  
 United Kingdom  
 E-mail: mdweck@staffmail.ed.ac.uk

<http://dx.doi.org/10.1016/j.jacc.2012.07.024>

## REFERENCES

1. Dweck MR, Chow MW, Joshi NV, et al. Coronary arterial  $^{18}\text{F}$ -sodium fluoride uptake: a novel marker of plaque biology. *J Am Coll Cardiol* 2012;59:1539-48.
2. Rogers IS, Nasir K, Figueroa AL, et al. Feasibility of FDG imaging of the coronary arteries: comparison between acute coronary syndrome and stable angina. *J Am Coll Cardiol* 2010;3:388-97.
3. Cheng VY, Slomka PJ, Le Meunier L, et al. Coronary arterial  $^{18}\text{F}$ -FDG uptake by fusion of PET and coronary CT angiography at sites of percutaneous stenting for acute myocardial infarction and stable coronary artery disease. *J Nucl Med* 2012;53:575-83.
4. Beheshti M, Saboury B, Mehta NN, et al. Detection and global quantification of cardiovascular molecular calcification by fluoro $^{18}$ -fluoride positron emission tomography/computed tomography—a novel concept. *Hell J Nucl Med* 2011;14:114-20.

## Efficiently Doing the Wrong Thing

We would like to commend Felker and Mentz (1) for their comprehensive review of volume removal therapy for patients with acute decompensated heart failure (ADHF). However, we also would like to challenge their most basic assertion that “Fluid retention and congestion are responsible for 90% of HF hospitalizations.” Neither of the references they cite (2,3) provide data to support that fluid retention occurs before the onset of ADHF in most patients. In fact, more recent studies that have measured weight gain (as a surrogate for fluid retention) show clearly that most patients gain either no, or minimal, weight before hospitalization for ADHF (4,5) despite increased cardiac filling pressures (5).

To account for increased cardiac filling pressures in the absence of weight gain, volume shifts rather than volume gains must occur. In a recent manuscript (6), we elucidate the likely mechanisms underlying these shifts, which involve a normally compliant splanchnic venous system that becomes noncompliant and results in redistribution of volume to the cardiopulmonary circulation, a process that can occur rapidly. We also point out that even in the minority of patients who do experience weight gain, little of the excess fluid resides within the effective circulatory volume (approximately 2.5%). The majority of patients presenting with congestion

likely have a syndrome of sympathetically mediated redistribution of volume and diuretic resistance.

Fluid retention and congestion do not necessarily represent the same phenomenon, and congestion most often is not due to fluid overload. Thus, strategies aimed at removing salt and water will necessarily be fraught with complications, including intravascular volume depletion, activation of the renin-angiotensin system, worsening renal function, and iatrogenically induced cardiorenal syndrome. Therefore, we caution readers to reconsider volume removal as the prime target of therapy for patients with ADHF. Diagnosing the contribution of redistribution versus total body salt and water gain emerges as the primary decision point, followed by therapy directed at the underlying etiology of congestion.

**\*Mark E. Dunlap, MD**  
**Paul A. Sobotka, MD**

\*MetroHealth Campus of Case Western Reserve University  
 Heart and Vascular Center  
 2500 MetroHealth Drive  
 Cleveland, Ohio 44109  
 E-mail: mdunlap@metrohealth.org

<http://dx.doi.org/10.1016/j.jacc.2012.07.025>

Please note: Dr. Dunlap is a consultant for Medtronic Inc., and has received research grants from Medtronic Inc./Ardian Inc., and BioControl Medical Inc. Dr. Sobotka is the Chief Medical officer for Coridea-NCI, and has been affiliated with Ardian, Inc., Medtronic, and CHF Solutions.

## REFERENCES

1. Felker GM, Mentz RJ. Diuretics and ultrafiltration in acute decompensated heart failure. *J Am Coll Cardiol* 2012;59:2145-53.
2. Lloyd-Jones D, Adams R, Carnethon M, et al. Heart disease and stroke statistics—2009 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 2009;119:e21-181.
3. Adams KF Jr., Fonarow GC, Emerman CL, et al. Characteristics and outcomes of patients hospitalized for heart failure in the United States: rationale, design, and preliminary observations from the first 100,000 cases in the Acute Decompensated Heart Failure National Registry (ADHERE). *Am Heart J* 2005;149:209-16.
4. Chaudhry SI, Wang Y, Concato J, Gill TM, Krumholz HM. Patterns of weight change preceding hospitalization for heart failure. *Circulation* 2007;116:1549-54.
5. Zile MR, Bennett TD, St John Sutton M, et al. Transition from chronic compensated to acute decompensated heart failure: pathophysiological insights obtained from continuous monitoring of intracardiac pressures. *Circulation* 2008;118:1433-41.
6. Fallick C, Sobotka PA, Dunlap ME. Sympathetically mediated changes in capacitance: redistribution of the venous reservoir as a cause of decompensation. *Circ Heart Fail* 2011;4:669-75.

## Reply

We thank Drs. Dunlap and Sobotka for their interest in our recent review on volume removal in patients with acute decompensated heart failure (ADHF). We agree with them that redistribution of volume (rather than increase in total body volume) may be an important and underappreciated mechanism in patients with acute decompensated heart failure, as our group has suggested previously (1).

Although we welcome “outside the box” thinking in a field that clearly would benefit from new ideas, we also suggest that the results from recent clinical trials might dampen the enthusiasm for

the hypothesis that redistribution of volume is the primary pathophysiology of most patients with ADHF. First, if redistribution were the primary culprit of decompensation, it would follow that vasodilator therapy would be a highly effective therapy for ADHF. Unfortunately, the largest vasodilator trial ever performed in heart failure, the recently completed ASCEND-HF (Acute Study of Clinical Effectiveness of Nesiritide in Decompensated Heart Failure) study, showed no significant clinical benefit of vasodilator therapy (2). Secondly, the randomized, double-blinded DOSE (Diuretic Optimization Strategies Evaluation) study suggested greater efficacy in terms of symptom relief and decongestion with a high-dose versus a low-dose loop diuretic regimen, without any evidence for long-term harm on renal function or clinical endpoints (3). These data would seem to contradict the assertion that excess volume is not an important mechanism for most patients with ADHF and that “strategies aimed at removing salt and water will necessarily be fraught with complications.”

Although both volume overload and volume redistribution are likely to be operative in individual patients to varying degrees, current diagnostic tools do not appear adequate to distinguish these mechanisms clinically. We strongly agree with the general suggestion in the final sentence of their letter that greater focus on the

specific mechanisms operative in individual patients could lead to better individualization of therapy and better outcomes in ADHF.

**\*G. Michael Felker, MD, MHS**  
**Robert J. Mentz, MD**

\*Duke Clinical Research Institute  
Division of Cardiology  
2400 Pratt Street  
Room 0311 Terrace Level  
Durham, North Carolina 27705  
E-mail: [michael.felker@duke.edu](mailto:michael.felker@duke.edu)

<http://dx.doi.org/10.1016/j.jacc.2012.07.023>

#### REFERENCES

1. Cotter G, Felker GM, Adams KF, Milo-Cotter O, O'Connor CM. The pathophysiology of acute heart failure—is it all about fluid accumulation? *Am Heart J* 2008;155:9–18.
2. O'Connor CM, Starling RC, Hernandez AF, et al. Effect of nesiritide in patients with acute decompensated heart failure. *N Engl J Med* 2011;365:32–43.
3. Felker GM, Lee KL, Bull DA et al. Diuretic strategies in patients with acute decompensated heart failure. *N Engl J Med* 2011;364:797–805.